

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY



0410

ANIGANE

ACTIVE CONSTITUENTS:

| | |
|--|----------|
| Sodium Glycerophosphate | 16.3 g/L |
| Potassium (as Potassium Citrate) | 9.1 g/L |
| Iron (as Ammonium Iron (III) Citrate) | 4.9 g/L |
| Manganese (as Manganese (II) Sulphate) | 0.8 g/L |

Increases liveweight gain and improves feed conversion in broilers. Acts as an appetite stimulant in pigs and poultry.

1 litre



014F

DIRECTIONS FOR USE:

PRECAUTION: LAYERS AND BREEDERS should not receive drinking water medicated with Anigane for more than 48 consecutive hours. Where loss of appetite is a continuing problem (e.g. during periods of heat stress) a once a week regime is suggested with a maximum of two days medication in every seven days.

DOSAGE AND ADMINISTRATION

SHAKE BEFORE USE

BROILERS: To increase liveweight gain and improve feed conversion, Anigane should be fed continuously from day old to market.

Anigane should be added to the drinking water in continuous flow-through systems. Add 1 litre of Anigane to each 1500 litres of drinking water.

Where **Automatic Water Medicators** are used adjust the unit to deliver a constant ration of 1:1500.

Note: When environmental temperatures exceed 30°C, or farm records show that at certain times of the year birds double their water intake, the concentration of Anigane should be adjusted to 1 litre of Anigane to 3000 litres of water.

In **Static Watering Systems** use as follows:

| Tank Size | Anigane Required |
|-----------|------------------|
| 5 litre | 4 mL |
| 25 litre | 17 mL |
| 50 litre | 35 mL |
| 100 litre | 68 mL |
| 200 litre | 135 mL |

Mix thoroughly.

BROILERS - 6 WEEKS OR OLDER: To reduce the effect of heat stress, Add 100 mL per 1000 broilers in sufficient drinking water which will be consumed in half a day.

Recommended dosings are late evening and early morning.

OTHER POULTRY (Layers, Duck, Turkeys): For appetite stimulation. Add 70 mL of Anigane for every 100 litres of drinking water consumed.

Anigane may be used in conjunction with other poultry medications without affecting potency.

As an appetite stimulant Anigane may be added to feed, milk or drinking water. It should be fed each day for at least five days.

PIGS

| | |
|-------------------|-------------------------------|
| Starters | Twice Daily 15 mL to 30 mL |
| Weaners & Growers | 20 mL to 30 mL |
| Breeders | 30 mL to 40 mL |

WITHOLDING PERIODS: 0 days

TRADE ADVICE

EXPORT SLAUGHTER INTERVAL (ESI): 0 days

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

CONDITIONS OF SALE: As the use of this product is beyond the control of the manufacturer, the purchaser assumes responsibility for any results which may occur and shall not hold the manufacturer liable for any loss or injury however occurring.

MANUFACTURED BY:

RUTH CONSOLIDATED INDUSTRIES PTY LTD

ABN 18 001 840 080

Unit 4, 26 Kent Road, Mascot NSW 2020 AUSTRALIA

Phone: 61 2 9667 0700

Fax: 61 2 9669 0430

Free Call: 1800 257 193

E-Mail: info@rci.com.au

Web Site: www.rci.com.au

DISPOSAL: Triple or (preferably) pressure rinse into the medicated water. Do not dispose of undiluted chemicals on-site. Return clean container to recycle point.

STORE BELOW 25°C (Air Conditioning)

APVMA Approval Number: 41414/1L/0410

Batch No:

Mfg. Date:

Exp. Date:



014B
1018