

FOR ANIMAL TREATMENT ONLY

SELECTOLYTE

ACTIVE CONSTITUENTS

SODIUM	109 g/kg
POTASSIUM	241 g/kg
BICARBONATE	289 g/kg

Selectolyte is a balanced source of essential body salts in a special form to counteract the effects of dehydration in all livestock

DIRECTIONS FOR USE:

Selectolyte may be used in drinking water, in feed, or as a drench.
Use can be continued until the condition is corrected.

POULTRY DOSE RATE:

To prevent dehydration, and weight loss during transportation:
Use at the rate of 1kg of Selectolyte in each 500 litres of drinking water.
Medicate the water for 5-10 days.

In cases of mild dehydration:

Use at the rate of 1kg of Selectolyte in each 150 litres of drinking water.
Medicate the water for 5-10 days.

In cases of severe dehydration:

Use at the rate of 1kg of Selectolyte in each 100 litres of drinking water.
Medicate the water for 5 days at this rate and then continue at the mild dehydration rate.

IMPORTANT: if adding to the drinking water supply always mix the required amount of Selectolyte with a small quantity of water first and stir thoroughly. Then add the resulting mixture to the water tanks and again stir thoroughly.

PIGS, CALVES AND DAIRY CATTLE DOSE RATE:

- a) In feed 2.5 kg per 1000 kg for 5-7 days.
- b) In water 0.5 kg per 100 litres drinking water for 5-7 days.
- c) As drench 10 kg per 100 kg of body weight daily for 5-7 days.

EXPORT SLAUGHTER INTERVAL (ESI): Zero (0) days.

WITHHOLDING PERIOD: NIL

MANUFACTURED BY: RUTH CONSOLIDATED INDUSTRIES PTY LTD

A.C.N. 001 840 080

Unit 5, 7-9 Kent Road, Mascot, NSW 2020, Australia

Phone: 61 2 9667 0700

Fax: 61 2 9669 0430

Free Call: 1800 257 193

E-mail: info@rci.com.au

Internet: www.rci.com.au

FIRST AID: If poisoning occurs contact a doctor or **Poisons Information Centre, Australia 13 11 26.**

DISPOSAL: Dispose of empty container by wrapping with paper and putting in garbage.

STORE BELOW 30°C (Room Temperature)

BATCH NO:

MFG DATE:

EXPIRY DATE:



RLP
Approved



25kg

APVMA No: 41416/ 50112